

Abstract Submitted to the
International Conference on Strongly Correlated Electron Systems
University of Michigan, Ann Arbor
August 6-10, 2001

Optical Transitions in Non-Stoichiometric LaMnO_3 Identifying Charge Transfer Transitions*

N.N. Kovaleva¹, J. Gavartin², A.V. Boris¹, and A. M. Stoneham²

¹ *Institute of Solid State Physics, Russian Academy of Sciences, Chernogolovka,
Moscow distr., 142432, RUSSIA*

² *University College London, Gower Street, London WC1E 6BT, UK*

Self-trapping of charge carriers is always associated with photo-induced charge transfer (CT) transitions. Using the Mott-Littleton approach we calculate energies of the main CT transitions involving Mn^{4+} and O^- localized holes, which determine specific electronic transport properties of doped CMR materials of $\text{La}_{(1-x)}\text{A}_x\text{MnO}_3$ (where A - divalent alkaline-earth ions). In these calculations extensive use is made of results on *in-crystal* ionization potentials of Mn^{3+} and O^{2-} ions from a companion paper*. Our Mott-Littleton calculations predict energies for the main optical bands, which match these observed in the experimental optical conductivity spectrum near 2.3, 5 and 9 eV in the LaMnO_3 crystal. Experimental low-energy optical conductivity bands well agree with our calculated CT transitions involving Mn^{4+} and O^- localized holes. The reasonable agreement with experiment of our predicted energies, linewidth and oscillator strength leads us to plausible assignment of the optical bands observed.

* N.N. Kovaleva, J. Gavartin, A. Shluger, A.V. Boris, and A. M. Stoneham, "Formation and Relaxation Energies of Electronic Holes in LaMnO_3 Crystal"

*This work was supported by the Royal Society/NATO